

Övriga referenser som ingått i depressionsprogrammet

Arroll B, Elley CR, Fishman T, Goodyear-Smith FA, Kenealy T, Blashki G, Kerse N, Macgillivray S: Antidepressants versus placebo for depression in primary care. *Cochrane Database Syst Rev.* 2009 (3):Cd007954. <http://www.ncbi.nlm.nih.gov/pubmed/19588448>

Barbato A, D'Avanzo B: Marital therapy for depression. *Cochrane Database Syst Rev.* 2006 (2):Cd004188. <http://www.ncbi.nlm.nih.gov/pubmed/16625597>

Barbui C, Cipriani A, Patel V, Ayuso-Mateos JL, van Ommeren M: Efficacy of antidepressants and benzodiazepines in minor depression: systematic review and meta-analysis. *Br J Psychiatry.* 2011 Jan;198(1):11-6, sup 1. <http://www.ncbi.nlm.nih.gov/pubmed/21200071>

Candy M, Jones L, Williams R, Tookman A, King M: Psychostimulants for depression. *Cochrane Database Syst Rev.* 2008 Apr 16;(2):CD006722. doi: 10.1002/14651858.CD006722.pub2.

Cipriani A, Koesters M, Furukawa TA, Nose M, Purgato M, Omori IM, Trespido C, Barbui C: Duloxetine versus other anti-depressive agents for depression. *Cochrane Database Syst Rev.* 2012;10:Cd006533. <http://www.ncbi.nlm.nih.gov/pubmed/23076926>

Cipriani A, Purgato M, Furukawa TA, Trespido C, Imperadore G, Signoretti A, Churchill R, Watanabe N, Barbui C: Citalopram versus other anti-depressive agents for depression. *Cochrane Database Syst Rev.* 2012 Jul 11;7:CD006534. doi: 10.1002/14651858.CD006534.pub2

Cipriani A, Santilli C, Furukawa TA, Signoretti A, Nakagawa A, McGuire H, Churchill R, Barbui C: Escitalopram versus other antidepressive agents for depression. *Cochrane Database Syst Rev.* 2009 (2):Cd006532. <http://www.ncbi.nlm.nih.gov/pubmed/19370639>

Cipriani A, La Ferla T, Furukawa TA, Signoretti A, Nakagawa A, Churchill R, McGuire H, Barbui C: Sertraline versus other antidepressive agents for depression. *Cochrane Database Syst Rev.* 2009 (2):Cd006117. <http://www.ncbi.nlm.nih.gov/pubmed/19370626>

Cipriani A, Brambilla P, Furukawa T, Geddes J, Gregis M, Hotopf M, Malvini L, Barbui C: Fluoxetine versus other types of pharmacotherapy for depression. *Cochrane Database Syst Rev.* 2005 (4):Cd004185. <http://www.ncbi.nlm.nih.gov/pubmed/16235353>

Cooney G, Dwan K, Mead G: Exercise for depression. *JAMA.* 2014 Jun 18;311(23):2432-3. doi: 10.1001/jama.2014.4930.

Cuijpers P, van Straten A, Schuurmans J, van Oppen P, Hollon SD, Andersson G: Psychotherapy for chronic major depression and dysthymia: a meta-analysis. *Clin Psychol Rev.* 2010 Feb;30(1):51-62. doi: 10.1016/j.cpr.2009.09.003.

Dennis CL, Hodnett E: Psychosocial and psychological interventions for treating postpartum depression. *Cochrane Database Syst Rev.* 2007 Oct 17;(4):CD006116.

Goss AJ, Kaser M, Costafreda SG, Sahakian BJ, Fu CH: Modafinil augmentation therapy in unipolar and bipolar depression: a systematic review and meta-analysis of randomized controlled trials. *J Clin Psychiatry.* 2013 Nov;74(11):1101-7. doi: 10.4088/JCP.13r08560.

Guaiana G, Barbui C, Hotopf M: Amitriptyline for depression. *Cochrane Database Syst Rev.* 2007 (3):Cd004186. <http://www.ncbi.nlm.nih.gov/pubmed/17636748>

Henken HT, Huibers MJ, Churchill R, Restifo K, Roelofs J: Family therapy for depression. *Cochrane Database Syst Rev.* 2007 (3):Cd006728. <http://www.ncbi.nlm.nih.gov/pubmed/17636850>

Huntley AL, Araya R, Salisbury C: Group psychological therapies for depression in the community: systematic review and meta-analysis. *Br J Psychiatry.* 2012 Mar;200(3):184-90.

Komossa K, Depping AM, Gaudchau A, Kissling W, Leucht S: Second-generation antipsychotics for major depressive disorder and dysthymia. *Cochrane Database Syst Rev.* 2010 Dec 8;(12):CD008121. doi: 10.1002/14651858.CD008121.pub2.

Leucht C, Huhn M, Leucht S: Amitriptyline versus placebo for major depressive disorder. *Cochrane Database Syst Rev.* 2012 Dec 12;12:CD009138. doi: 10.1002/14651858.CD009138.pub2.

Magni LR, Purgato M, Gastaldon C, Papola D, Furukawa TA, Cipriani A, Barbui C: Fluoxetine versus other types of pharmacotherapy for depression. *Cochrane Database Syst Rev.* 2013;7:Cd004185. <http://www.ncbi.nlm.nih.gov/pubmed/24353997>

Omori IM, Watanabe N, Nakagawa A, Cipriani A, Barbui C, McGuire H, Churchill R, Furukawa TA: Fluvoxamine versus other anti-depressive agents for depression. *Cochrane Database Syst Rev.* 2010 (3):Cd006114. <http://www.ncbi.nlm.nih.gov/pubmed/20238342>

Rethorst CD, Trivedi MH: Evidence-based recommendations for the prescription of exercise for major depressive disorder. *J Psychiatr Pract.* 2013 May;19(3):204-12. doi: 10.1097/01.pra.0000430504.16952.3e.

Semkovska M1, McLoughlin DM: Objective cognitive performance associated with electroconvulsive therapy for depression: a systematic review and meta-analysis. *Biol Psychiatry.* 2010 Sep 15;68(6):568-77. doi: 10.1016/j.biopsych.2010.06.009. Epub 2010 Jul 31.

Socialstyrelsen. Nationella riktlinjer för vård vid depression och ångestsyndrom 2010.

Spielmans GI, Berman MI, Linardatos E, Rosenlicht NZ, Perry A, Tsai AC: Adjunctive atypical antipsychotic treatment for major depressive disorder: a meta-analysis of depression, quality of life, and safety outcomes. *PLoS Med.* 2013;10(3):e1001403. doi: 10.1371/journal.pmed.1001403. Epub 2013 Mar 12.

Taylor MJ, Wilder H, Bhagwagar Z, Geddes J: Inositol for depressive disorders. *Cochrane Database Syst Rev.* 2004;(2):CD004049

Wijkstra J, Lijmer J, Burger H, Geddes J, Nolen WA: Pharmacological treatment for psychotic depression. *Cochrane Database Syst Rev.* 2013 Nov 26;11:CD004044. doi: 10.1002/14651858.CD004044.pub3.